

Running head: EMOTIONAL PROCESSING IN PSYCHOPATHY VARIANTS

Primary and Secondary Variants of Juvenile Psychopathy Differ in
Emotional Processing

Eva R. Kimonis¹, Paul J. Frick², Elizabeth Cauffman³, Asha Goldweber⁴, & Jennifer Skeem³

Author Note

¹University of South Florida, ²University of New Orleans, ³University of California, Irvine,

⁴Johns Hopkins University

Funding for this study was provided to Elizabeth Cauffman, Ph.D. from the National Institute of Mental Health (K01MH01791-01A1) and from the Center for Evidence-Based Corrections. We are especially grateful to the many individuals responsible for the data collection and preparation.

Address correspondence to Eva R. Kimonis, Ph.D., Assistant Professor, Louis de la Parte Florida Mental Health Institute, Mental Health Law and Policy, University of South Florida, 13301 Bruce B. Downs Blvd. MHC 2639, Tampa, FL 33612, (813) 974-8612 Office (813) 974-6411 Fax, Email: ekimonis@usf.edu.

Manuscript submitted for a Special Issue of *Development and Psychopathology* on Multi-level Approaches toward Understanding Antisocial Behavior: Current Research and Future Directions

Abstract

Accumulating research suggests that psychopathy can be disaggregated into low-anxious primary and high-anxious secondary variants and this research may be important for understanding antisocial youths with callous-unemotional traits. Using model-based cluster analysis, the present study disaggregated 165 serious male adolescent offenders (M age= 16) with high scores on the *Youth Psychopathic Traits Inventory* (Andershed, Kerr, Stattin, & Levander, 2002) into primary and secondary variants based on the presence of anxiety. Results indicated that the secondary, high-anxious variant was more likely to show a history of abuse and scored higher on measures of emotional and attentional problems. On a picture version of the dot probe task, the low-anxious primary variant was not engaged by emotionally distressing pictures, whereas the high-anxious secondary variant was more attentive to such stimuli (Cohen's $d = .71$). Although the two groups differed as hypothesized from one another, neither differed significantly in their emotional processing from a nonpsychopathic control group of offending youth ($n = 208$). These results are consistent with the possibility that the two variants of psychopathy, both of which were high on callous-unemotional traits, may have different etiological pathways, with the primary being more related to a deficit in the processing of distress cues in others and the secondary being more related to histories of abuse and emotional problems.

Keywords: juvenile psychopathy; primary and secondary psychopathy; emotional processing; juvenile offenders

Primary and Secondary Variants of Juvenile Psychopathy Differ in Emotional Processing

Research on the development of criminal behavior has long recognized the great heterogeneity within youth who show serious aggressive and antisocial behaviors (Moffitt, 1993). That is, within youth diagnosed with Conduct Disorder or within youth who show serious criminal behavior, there appears to be distinct groups that differ on the life course of their antisocial behavior, the severity of their antisocial behavior, the likely causal process leading to their antisocial behavior, and the most effective treatment for their antisocial behavior (Frick & Viding, 2009). As a result, there has been great interest in defining methods for distinguishing among the different subgroups of antisocial individuals. One particularly promising method that is being considered for inclusion in the upcoming 5th revision of the Diagnostic Statistical Manual of Mental Disorders is distinguishing between those who do and do not show significant callous-unemotional (CU) traits (Frick & Moffitt, 2010).

CU traits are characterized by a lack of empathy or guilt, and callous use of others (Frick, 2009). These traits have often been considered key features to the broader construct of psychopathy (Cleckley, 1941; Cooke & Michie, 2001; Hare, 2003). Youth high on CU traits represent only a minority of antisocial youths in either community, clinic-referred, or forensic samples (Frick & Moffitt, 2010; Kahn, Frick, Youngstrom, Findling, & Youngstrom, in press; Kruh, Frick, & Clements, 2005; Rowe, et al., 2010). Yet, this group appears to be significantly different from other antisocial youths in a number of ways that cut across multiple levels of analyses.

Specifically, youths with CU traits differ behaviorally by showing a more stable pattern of behavior that is characterized by more conduct problems and aggression, especially instrumental aggression (Frick, et al., 2003; Frick, Stickle, Dandreaux, Farrell, & Kimonis,

2005). They also differ on the relative strength of genetic and environmental influences on their behavior (Fontaine, Rijdsdijk, McCrory, & Viding, 2010; Viding, Blair, Moffitt, & Plomin, 2005; Viding, Jones, Frick, Moffitt, & Plomin, 2008). For example, Viding et al. (2005) demonstrated that the heritability of antisocial behavior at age 7 for those high on CU traits was considerably greater ($h=.81$) than for those low on CU traits ($h=.30$). Further, several studies have shown that the antisocial behavior of youths with CU traits are less strongly related to problematic parenting factors (Edens, Skopp, & Cahill, 2008; Hipwell, et al., 2007; Oxford, Cavell, & Hughes, 2003; Wootton, Frick, Shelton, & Silverthorn, 1997).

In addition to these differences in the genetic and environmental influences, youth scoring high on measures of CU traits differ from other youth with conduct problems on various physiological measures as well, including evincing lower resting salivary cortisol (i.e., stress hormone) levels (Loney, Butler, Lima, Counts, & Eckel, 2006) and reduced amygdala activation while processing fearful expressions (Jones, Laurens, Herba, Barker, & Viding, 2009; Marsh, et al., 2008). They also show cognitive differences, such as decreased sensitivity to punishment and threat cues and a tendency to overestimate the potential positive consequences of acting aggressively compared to those with lower levels of CU traits (Blair, Peschardt, Budhani, Mitchell, & Pine, 2006; Fisher & Blair, 1998; Pardini, Lochman, & Frick, 2003). Finally, youth high on CU traits show differences in emotional processing compared to other antisocial youth. For example, these youth are more likely to show impaired recognition of and reduced sensitivity to negative emotional stimuli, especially stimuli that typically evoke the experience of empathy (e.g., pictures, words or sounds depicting others' distress; Blair, 1999; Blair, Colledge, Murray, & Mitchell, 2001; Dadds, El Masry, Wimalaweera, & Guastella, 2008; Kimonis, Frick, Fazekas, & Loney, 2006; Kimonis, Frick, Muñoz, & Aucoin, 2008).

Thus, antisocial youths with and without CU traits seem to show very different behavioral, genetic, contextual, cognitive, and emotional characteristics. As a result of these differences, any comprehensive causal theory of antisocial behavior must explain these differences in youths with and without CU traits (Frick, Blair, & Castellanos, in press; Frick & Viding, 2009). Further, it is also important to derive theories for how children may develop these traits. Many theories have linked CU traits to a fearless temperament (Frick, 2006; Lykken, 1995; Patrick, Fowles, & Krueger, 2009). This temperament is thought to leave a child insensitive to emotional cues (e.g., angry parent signaling punishment, crying peer signaling distress) that normally contribute to the development of empathy, guilt, and other aspects of the affective components of conscience (Fowles & Kochanska, 2000; Kochanska, 1993).

As noted above, CU traits are a critical dimension included in most definitions of psychopathy. However, CU traits are only one facet of psychopathy, often termed the “affective discomfort” component (Hare & Neumann, 2010). More importantly, some authors have suggested that the CU traits component of psychopathy may designate an etiologically distinct subtype of psychopathy. For example, Lykken (1957; 1995) contrasts what has come to be called the primary psychopath—described by Cleckley (1955) as having a core deficit in emotional responding—against the impulsive secondary psychopath who is prone to negative emotionality and neuroticism. This theoretical model would suggest that persons with primary psychopathy manifest marked CU traits, whereas those with the secondary variant would manifest greater impulsivity, hostility, and behavioral deviance. There has been only mixed support for this distinction. That is, secondary variants often are shown to show higher rates of impulsivity and negative emotionality; however, both variants tend to show higher rates of CU traits (Hicks,

Markon, Patrick, Krueger, & Newman, 2004; Vassileva, Kosson, Abramowitz, & Conrod, 2005) (cf. Poythress, et al., 2010; Skeem, Johansson, Andershed, Kerr, & Louden, 2007).

As a result of these findings, others theories to distinguish between the two variants have focused more on the presence of anxiety (Karpman, 1941). Although major theories of psychopathy often feature emotional deficits like fearlessness or a lack of anxiety (Cleckley, 1941), such traits are not included in leading psychopathy measures (Forth, Kosson, & Hare, 2003; Hare, 2003; Hart, Cox, & Hare, 1995) and individuals labeled as psychopathic can vary substantially in their levels of anxiety. More importantly, those individuals with psychopathic traits who are high and low on measures of anxiety seem to show differences in their emotional and cognitive processing. For example, Newman and colleagues reported that adult psychopathic offenders with high scores on measures of anxiety failed to show the same deficits in passive avoidance learning, modulation of responses to emotional and neutral stimuli, and fear-potentiated startle response as their psychopathic counterparts with low anxiety (Newman, Patterson, Howland, & Nichols, 1990; Newman, Schmitt, & Voss, 1997; Newman & Schmitt, 1998).

These findings by Newman and colleagues would be consistent with early research that distinguished variants of psychopathy focusing largely on the presence of anxiety.¹ For example, Karpman's description of a person with secondary psychopathy focused largely on the presence of high levels of anxiety (Karpman, 1948b, p. 523). Further, Karpman viewed secondary psychopathy as an acquired disturbance in emotional functioning stemming chiefly from abusive parenting or other childhood trauma (Karpman, 1941; 1948a; 1948b; Skeem, Poythress, Edens, Lilienfeld, & Cale, 2003). Significant research has provided support for Karpman's distinction. That is, there appears to be a variant of psychopathy associated with higher levels of anxiety, as

well as greater histories of childhood abuse and trauma in both studies of *youth* with high scores on measures of psychopathy (Kimonis, Skeem, Cauffman, & Dmitrieva, 2011; Tatar, Cauffman, Kimonis, & Skeem, in press; Vaughn, Edens, Howard, & Smith, 2009), and *adults* with high scores on measures of psychopathy (Hicks, et al., 2004; Hicks, Vaidyanathan, & Patrick, 2010; Skeem, et al., 2007). Also, consistent with Karpman's conceptualization of secondary psychopathy as an acquired emotional disturbance, studies have shown that those with secondary psychopathy also manifest greater depression, attentional problems, and anger (Kimonis, et al., 2011; Lee, Salekin, & Iselin, 2010; Vaughn, et al., 2009).

Thus, research has largely supported the presence of two variants of psychopathy in samples of both adults and adolescents, both of which tend to be high on CU traits. One significant limitation to this research is a failure to compare these different groups on their emotional processing. As noted earlier, deficits in the processing of distress cues in others have played a key role in theories trying to explain the development of CU traits. Further, theories to explain the development of primary psychopathy have often focused on a dispositional deficit in the person's emotional responsiveness to others, with persons with primary psychopathy showing less emotional responsiveness to negative emotional cues which makes them less sensitive to distress cues in others and to cues for punishment (Cleckley, 1976).

In contrast, the experience of maltreatment and related trauma that is characteristic of secondary psychopathy has been linked with heightened emotional sensitivity to negative emotional stimuli. For example, physically abused children show heightened emotional reactivity and hypervigilance to threat-related cues (Dodge & Pettit, 2003). Similarly, Pollak and colleagues reported that maltreated children showed abnormalities in their processing of angry faces and their regulation of emotional states (Pollak & Kistler, 2002; Pollak, Klorman,

Thatcher, & Cicchetti, 2001; Pollak & Sinha, 2002; Pollak & Tolley-Schell, 2003). For example, children who were physically abused tended to over-attend to angry expressions and experienced difficulty disengaging from such cues, even when they were task-irrelevant. Pollak and colleagues suggest that this pattern of emotional processing is likely an adaptive mechanism for maltreated youth who must rapidly filter out and identify anger cues that signal impending threat.

Thus, persons with primary and secondary psychopathy may have very different patterns of responses to emotional stimuli, both of which could contribute to their problems in conscience development. That is, theories of moral development suggest that patterns of emotional underarousal, on one hand, and overarousal, on the other, may both interfere with the normative development of conscience (Frick & Morris, 2004; Hoffman, 1982; Young, Fox, & Zahn-Waxler, 1999). Unfortunately, very little research has tested this possibility. In one notable exception, Bagley et al. (2009) found that adult males with primary psychopathy made significantly more errors in recognizing emotional tones conveyed in sentences spoken in a foreign language (i.e., prosodic condition) relative to nonpsychopaths, whereas those with secondary psychopathy did not differ significantly from either group. However, they also found that the primary and secondary variants both made significantly more errors than nonpsychopaths in recognizing affective cues conveyed in sentences spoken in a neutral and flat tone of voice (i.e., semantic condition: content of language), but did not differ significantly from each other. Importantly, although not significant, the primary group made 13% more errors relative to nonpsychopaths, whereas the secondary group made 9% more errors. This difference was most pronounced for sadness cues where the group difference between primary and secondary variants yielded a moderate effect size ($d = 0.41$). These findings suggest that primary and secondary psychopathy variants may show at least some differences in their processing of

emotional stimuli. It is possible that emotion tasks that rely less on effortful processing of emotional information (i.e., encoding sentences) and more on automatic processing, such as the dot probe task described below, might be more sensitive to differences among psychopathy variants.

The Present Study

In summary, the presence of CU traits has been important for designating a distinct group of antisocial youths who differ behaviorally, neurologically, socially, cognitively, and emotionally from antisocial youths without these traits. Importantly, CU traits are key features to most conceptualizations of psychopathy. Thus, the decades of research on psychopathic traits could inform theories on the development of CU traits. In particular, research suggests that there may be primary and secondary variants of psychopathy. Some authors have suggested that CU traits may be more specific to the primary variant but this research has not been conclusive. What has been shown more consistently in past research is that the secondary variant shows higher rates of anxiety, has more extensive histories of abuse, and shows higher rates of emotional and attentional problems. However, a key limitation to this research is the lack of research testing whether or not the primary group is more likely to display deficits in the automatic processing of distress cues in others.

Thus, to advance this research, we tested whether distinct groups of juvenile offenders with high rates of psychopathic traits could be identified using levels of anxiety to form this typology. Based on past research, we predicted that two groups of youths high on psychopathy would emerge, with one group showing high rates of anxiety (i.e., a secondary group) and the other (i.e., a primary group) showing low to average levels of anxiety. Also, based on past research, we predicted that the group high on anxiety would be more likely to have histories of

abuse and would show higher rates of emotional and attentional problems. We further tested whether the two groups would differ in their level of CU traits but, based on past research, we did not predict differences on this variable. This test was still important, however, because if both groups were high on CU traits, then the primary and secondary distinction would be critical for understanding youth high on these traits. However, our biggest advance from past work is that we tested the hypothesis that the group low on anxiety, but not the group high on anxiety, would show deficits in the processing of emotional stimuli depicting distress in others.

Method

Participants

The data for the present study come from assessments of 373 male juvenile offenders housed in a secure confinement facility in Southern California. All youth between the ages of 14 and 17 years who were newly admitted to the facility, or returning on a new offense, were eligible to enroll in the study. The majority (94%; $n=350$) of youth (M age = 16.42, $SD = .79$; see Table 1) were from ethnic minority backgrounds (53% Hispanic, 29% African American, and 12% other [bi- or multi-racial]; 6% Caucasian), an ethnic composition that is representative of youth incarcerated in this region of the US (Snyder & Sickmund, 2006). The majority of study participants were sentenced for violent (e.g., murder, rape, robbery, aggravated assault) committing offenses (69%, $n=258$).

[Insert Table 1]

Procedure

All study procedures were approved by a University Institutional Review Board and a Certificate of Confidentiality was secured from the Department of Health and Human Services to ensure that the information disclosed by the youths remained confidential. Ninety seven percent

of parents/guardians contacted consented to their child's participation in the study; the youth assent rate was 96%. Youth completed a two-hour baseline interview within 48 hours of arrival to the facility as well as weekly and monthly follow-up interviews (i.e., weeks 2, 3, 4, and month 2). Each interview consisted of a battery of developmental, behavioral, emotional, and attitudinal measures.

Measures

Participants self-reported their age, ethnicity, and whether they had been to the facility before. Measures described below are categorized into clustering measures used in model-based cluster analysis (MBC) to identify primary and secondary psychopathy variants, and measures used to support the external validity of the resultant clusters based on theoretical conceptualizations of psychopathy variants.

Clustering measures.

Psychopathy. The Youth Psychopathic Traits Inventory (YPI; Andershed et al., 2002) is a 50-item self-report measure of psychopathic traits that was administered at the week 3 assessment. The YPI was developed in a community-based sample of youth with an aim to capture core affective and interpersonal personality features of psychopathy to the exclusion of the “more behavioral consequences of psychopathic personality traits” (Andershed et al., 2002, p. 135), or antisocial deviance and criminal behavior that some believe develop as a result of such traits (Skeem & Cooke, 2010a; 2010b) (cf. Hare & Neumann, 2010). The items of the YPI were written to assess ten target traits/scales in a relatively comprehensive (5 items per trait) and indirect, nontransparent manner (e.g., “I usually feel calm when other people are scared”). Prior factor analytic research supports a three-factor structure in which each higher-order factor is composed of several lower-order scales (in parentheses): Grandiose-Manipulative (i.e.,

interpersonal traits: dishonest charm, grandiosity, lying, manipulation), Callous-Unemotional (affective traits: remorselessness, callousness, unemotionality), and Impulsive-Irresponsible (lifestyle traits: impulsiveness, irresponsibility, thrill seeking) (Andershed et al., 2002). The YPI correlates moderately with the Psychopathy Checklist: Youth Version (PCL:YV, $r=.35$, Forth, et al., 2003), with low to moderate correlations with a variety of self-report conduct problem indices, supporting its convergent validity (Cauffman, Kimonis, Dmitrieva, & Monahan, 2009). In the current sample, the YPI total ($\alpha=.90$) and factor scores (.71 [affective] to .89 [interpersonal]) demonstrated adequate internal consistency.

Anxiety. The Revised Children's Manifest Anxiety Scale (RCMAS; Reynolds & Richmond, 1985; 2000) is a standardized measure designed to assess anxiety in youth between the ages of 6 and 19. The RCMAS has 37 items divided into four scales: Physiological Anxiety (10 items; e.g., "Often I feel sick in my stomach," $\alpha=.64$), Worry/Oversensitivity (11 items; e.g., "I worry about what is going to happen," $\alpha=.76$), Social Concerns/Concentration (7 items; "A lot of people are against me," $\alpha=.68$), and a Lie Scale (9 items; e.g., "I never get angry," not included in analyses). The child responds to each question with a "Yes" or "No" answer. The RCMAS possesses moderate test-retest reliability over a 9-month period ($r=.63$; Reynolds, 1981), and its construct validity is supported by past research (e.g., Muris, Merckelbach, Ollendick, King, & Bogie, 2002).

External criteria variables.

Maltreatment history. The Life Events Scale (Gil-Rivas, 2003) consists of a list of 32 stressful life events. The youth is asked to endorse which of these events he has ever experienced in his lifetime. Maltreatment history was computed by totaling the seven items tapping parental

absence, domestic violence exposure, parental neglect, and physical, emotional, and sexual (two items) abuse ($\alpha=.64$).

Depression. The Center for Epidemiologic Studies Depression Scale (CES-D; Radloff, 1977) is a 20-item self-report measure of depressed mood, lack of positive affect, somatic symptoms, and interpersonal difficulties. A total score was computed by summing all 20 items rated at baseline, with higher scores indicative of more depressive symptomatology ($\alpha=.82$).

Psychopathology. The 112-item Child Behavior Checklist Youth Self Report (CBCL-YSR; Achenbach, 1991) was used to assess negative emotionality/internalizing psychopathology. The withdrawn/depressed (7 items), anxious/depressed (16 items), and attention problems (9 items) subscales rated at baseline were used.

Anger. The Novaco Anger Scale (NAS; Novaco, 2003) is a 48-item self-report measure of various dimensions of anger. The total score rated at baseline ($\alpha=.90$) was used in the current study. The NAS is highly correlated with several other measures of anger, such as the STAXI Trait Anger Scale ($r=.84$; Novaco, 2003).

Emotional processing. The Emotional Pictures Dot-Probe task (Kimonis, Frick, et al., 2006) is a variant of the traditional word version of the task that has been used extensively in the anxiety literature (MacLeod, Mathews, & Tata, 1986). It is a spatially-oriented selective attention task that is administered via computer to capture automatic attentional bias toward emotional cues, providing an indirect index of emotional reactivity (see Schippell, Vasey, Cravens-Brown, & Bretveld, 2003). The version used in the current study was developed using primarily slides taken from the International Affective Picture System (IAPS; Lang, Bradley, & Cuthbert, 1997) that were carefully selected to tap distressing (e.g., crying child), positive (e.g., puppies), and neutral emotional content (e.g., fork). Because the number of neutral and distressing images was

not sufficient for dividing the slides into neutral, distress, and positive categories, additional slides (distress $n = 19$; neutral $n = 42$) were added that directly matched the IAPS slide content. For example, additional slides of a crying child were added to the existing IAPS slides of crying children (see Kimonis, et al., 2008). A recent meta-analysis suggests that selective attention is similarly robust for both pictorial and verbal stimuli (Bar-Haim, Lamy, Pergamin, Bakermans-Kranenberg, & van Ijzendoorn, 2007).

The dot probe task consists of one block of practice stimuli (16 picture pairs) followed by four test blocks of picture pairs, each containing 24 picture pairs. Each picture pair presentation consists of three sequential and non-overlapping components: (1) a 500 millisecond fixation cross appearing in the center of the screen, (2) a 250 millisecond simultaneous presentation of two picture stimuli that are centered and located immediately above and below the location of the fixation cross, and (3) an asterisk (i.e., dot probe) appearing in either the top or bottom picture location immediately after the offset of the picture. The objective of the task is to select a key on the keyboard that corresponds to the location on the screen (top or bottom) where the dot-probe appears, as quickly as possible. The time between when the probe appears and when the youth presses the corresponding key to its location is recorded in milliseconds and used for the calculation of facilitation indices (MacLeod & Mathews, 1988): $\text{Facilitation} = 1/2 \times [(\text{Neutral Only/Probe Top} - \text{Distress Up/Probe Top}) + (\text{Neutral Only/Probe Bottom} - \text{Distress Down/Probe Bottom})]$. The assumption is that if the spatial location of the probe corresponds to where the participant's attention is allocated, then their response to the probes' location will be faster than if their attention was allocated elsewhere. If no key is pressed within 5000 milliseconds, the response is recorded as incorrect. Consistent with past uses of the task (Vasey, Daleidon, Williams, & Brown, 1995; Vasey, El-Hag, & Daleidon, 1996), incorrect responses were not

included in the calculation of facilitation indices as they reflect that the participant was not paying attention to a specific stimulus pair. Higher scores reflect greater attentional orienting to emotional stimuli (see Kimonis, Cauffman, & Goldweber, 2009; Kimonis, Frick, et al., 2006; Kimonis, et al., 2008). The facilitation index to distress demonstrated adequate internal consistency in the current study ($\alpha=.81$). A subsample of youth in the psychopathic group ($n = 127$) completed the emotional pictures dot-probe task. This subsample did not differ significantly from those who did not complete the task ($n = 38$) on any of the study variables.

Results

Data Analyses

Descriptive statistics are provided in Table 1 on the clustering variables, anxiety and psychopathy. YPI and anxiety scores were uncorrelated with facilitation indices from the dot-probe task. The mean facilitation to distress pictures was negative ($M=-19.19$, $SD=56.30$ ms), in keeping with Kimonis et al. (2008) (see also Legerstee, et al., 2009). In incarcerated samples of youth, there may be no general facilitation effect for images of distress.

Consistent with prior research (Hicks, et al., 2004; Hicks, et al., 2010; Kimonis, Skeem, & Cauffman, 2006; Skeem, et al., 2007; Tatar, et al., in press), we first used model-based cluster analysis (MBC; Banfield & Raftery, 1993) to determine whether primary and secondary variants of juvenile psychopathy can be identified in a subsample ($n=165$) of male juvenile offenders scoring high on the measure of psychopathy. To validate the resultant clusters we used multivariate analyses of variance (MANOVA) and analyses of variance (ANOVA) to compare them on theoretically relevant factors not used to derive them. The nonpsychopathic comparison group was included in these analyses along with primary and secondary psychopathy variants.

Cluster Derivation

Although we recognize that the most compelling data suggest that psychopathy is a dimensional trait rather than categorical (Edens, Marcus, Lilienfeld, & Poythress, 2006; Lilienfeld, 1994; 1998; Marcus, Lilienfeld, Edens, & Poythress, 2006) (cf. Harris, Rice, & Quinsey, 1994; Vasey, Kotov, Frick, & Loney, 2005), we used a threshold score to select youth because our interest was in identifying subgroups within those with substantial psychopathic traits. Thus, to derive clusters of primary and secondary variants of psychopathy, those scoring greater than 121.5 on the YPI ($n=165$) were selected for analyses. Although the YPI does not have an established cut score for classifying youth as psychopathic, a score of 121.5 was found to correspond to a PCL:YV score of 30 in a large sample ($N=1,171$) of adolescent male offenders (Cauffman, et al., 2009). We also focused on youth with high scores on the YPI (rather than all youthful offenders) for the cluster analyses to permit comparison of the results with prior studies of adult (Hicks, et al., 2004; Skeem, et al., 2007) and juvenile variants of psychopathy (Kimonis, et al., 2011; Vaughn, et al., 2009) which used this methodology. The remaining youth ($n=208$) with YPI total scores of 121.5 and below were used as a comparison group. The characteristics of the general sample ($N=373$), clustering subsample ($n=165$) and comparison subsample ($n=208$) are provided in Table 1. Relative to the comparison group, the group scoring high on the YPI (i.e., clustering group) obtained significantly higher scores on the RCMAS, and was more likely to have a property-related committing offense.

MBC was performed using SPLUS 7.0 (Insightful Corporation, 1988-2005), the *mclust* library (Fraley & Raftery, 2002a; 2002b; 2008), and the expectation maximization (EM) algorithm. MBC reduces some of the uncertainties inherent in common clustering methods by testing the relative fit of ten models that vary in their assumptions about the structure of the data. More detailed discussions of MBC are provided by Hicks et al. (2004) and Skeem et al. (2007).

Youths' Z scores (based on the psychopathic subsample) on the three higher-order factors of the YPI and the three anxiety scales of the RCMAS were used as clustering variables. According to BIC values generated by MBC, models that specified one cluster fit relatively poorly, indicating there were subgroups of youth scoring high on psychopathy with distinctive trait patterns. The best-fitting model was a two-cluster solution (cluster 1, $n = 122$; cluster 2, $n = 43$) with equal shape, equal volume, and diagonal orientation (BIC = -5236). The average classification certainty, or posterior probability that an individual was correctly assigned to a cluster, was high at 99.9%. Three-quarters of the sample had a fairly high ($\geq 97.5\%$) probability of correct assignment to a cluster suggesting confidence in this clustering solution.²

Description of Clusters

Using t-tests, the two clusters (C1, $n = 122$; C2, $n = 43$) were compared on total, factor, and subscale scores of the YPI and the RCMAS for descriptive purposes (see Figure 1). The second cluster ($n = 43$), which was labeled 'secondary,' reported significantly greater anxiety (physiological [$t(163)=-8.59$, $d = 1.35$]; worry [$t(163)=-13.61$, $d = 2.13$]; and social concerns/concentration [$t(163)=-11.92$, $d = 1.87$]; all p 's $<.001$), higher total YPI ($t(163)=-3.31$, $d = .52$, $p<.001$) and Impulsive-Irresponsible lifestyle factor scores ($t(163)=-3.70$, $d = .58$, $p<.001$), but not Callous-Unemotional or Grandiose-Manipulative scores, compared with the first cluster ($n = 122$) that was labeled 'primary.' Comparison youth were slightly more anxious than primary variants but significantly less anxious than secondary variants.

Validating Clusters: Maltreatment and Negative Emotionality/Symptomatology

By definition, clusters will differ on the variables used to cluster them. Thus, clusters were validated by comparing them via MANOVA on theoretically-relevant dimensions not used to derive them and found in prior empirical research to distinguish psychopathy variants.

Consistent with expectations, the secondary variant group reported significantly more maltreatment ($F(2,371) = 5.64, p < .01, d = .41$) compared with primary variants and comparison youth (see Table 2).³ Secondary variants also scored significantly higher on each measure of negative emotionality/symptoms compared with primary variants and comparison youth (Wilks' Lambda = 0.82, $F(10,732) = 7.68, p < .001, d = .65$). As depicted in Table 2, secondary variants endorsed significantly greater symptoms of depression/ social withdrawal and attention problems compared with primary variants and nonpsychopathic comparison youth. Also, relative to other groups, secondary variants endorsed significantly greater anger problems; primary variants also reported significantly more anger than nonpsychopathic comparison youth.

[Insert Table 2]

Validating Clusters: Processing of Distressing Emotional Stimuli

The two psychopathy clusters and comparison youth were also compared on their processing of distressing and positive emotional stimuli. Results of one-way ANOVAs revealed a significant but modest difference among the three groups in attentional orienting to distressing ($F(2,282) = 4.61, p < .05, d = .36$), but not positive stimuli. Given a priori hypotheses, planned comparisons between primary and secondary variants were conducted. Analyses revealed that primary and secondary variants differed significantly from each other (unequal variances; $t(43) = 2.34; p < .05$, Cohen's $d = .71$), but not from comparison youth (P v. C, unequal variances; $t(220) = -1.65; p = n.s.$, Cohen's $d = .22$; S v. C, unequal variances; $t(41) = 1.61; p = n.s.$, Cohen's $d = .50$; see Figure 2). That is, primary variants were not attentionally engaged by distressing stimuli, whereas secondary variants were more attentive to stimuli depicting distressing emotional content.

[Insert Figure 2]

Discussion

The results of this study are consistent with past research indicating that the construct of psychopathy is heterogeneous (Poythress & Skeem, 2005; Skeem, et al., 2003) and that high-anxious secondary variants endorse more negative emotionality and childhood abuse than low-anxious primary variants (Hicks, et al., 2004; Hicks, et al., 2010; Skeem, et al., 2007; Tatar, et al., in press; Vaughn, et al., 2009). The primary contribution of these results, however, is demonstrating that the variants of juvenile psychopathy—identified via cluster analysis—differed in their processing of emotional stimuli. As hypothesized, we found that secondary psychopaths were more engaged by distressing emotional stimuli than primary psychopaths (Cohen's $d = .71$).

Consistent with our findings, Bagley et al. (2009) also found that adult secondary psychopathy variants made fewer errors in recognizing the emotional tone and in recognizing sad emotional content ($d = 0.41$) of spoken sentences, relative to primary variants, albeit nonsignificantly. Our findings are also consistent with a body of research generated by Newman and colleagues finding that high-anxious adult psychopathic offenders—most closely resembling secondary variants—do not show deficits in passive avoidance learning, modulation of responses to emotional and neutral stimuli, and fear-potentiated startle response that are found to differentiate low-anxious psychopathic individuals (Newman, et al., 1990; Newman, et al., 1997; Newman & Schmitt, 1998). Together, these studies provide some support for our hypothesis that low-anxious primary variants, but not high-anxious secondary variants, will show emotional deficits believed by some to be core to the psychopathic personality (Cleckley, 1941).

Importantly, our findings are consistent with past research suggesting that the two groups are *not* distinguished by differences in their level of CU traits (Hicks, et al., 2004; Vassileva, et

al., 2005). This finding is important because, as noted previously, CU traits have been increasingly used to designate an important subgroup of antisocial youths. These findings suggest that those high on CU traits could fall into either the primary or secondary groups. Thus, causal theories of CU traits need to account for this heterogeneity.

Although this cross-sectional study cannot conclusively address etiological issues, we use our results to offer two speculative hypotheses that could be addressed in future research. One possibility is that these differences in emotional processing signal differences in temperament. Theories of moral development suggest that both “too high” and “too low” levels of emotional reactivity can impair conscience development and related complex social emotions of guilt and empathy (Kochanska, 1993; 1995; 1997). Whereas children with a fearless temperament (i.e., primary variants) may be insufficiently engaged by important socializing cues, highly emotionally reactive/ dysregulated children (i.e., secondary variants) might miss them because they are easily overwhelmed in negatively-charged situations where such cues tend to be elicited (e.g., parental anger, peer distress; see Frick & Morris, 2004; Hoffman, 1982; Young, Fox, & Zahn-Waxler, 1999). In short, our results would be consistent with the possibility that the two variants have different temperaments that are linked to CU traits and subsequent antisocial behavior via different emotional mechanisms.

Another possibility that would be consistent with our findings is that these differences in emotional processing signal differences in experience. Specifically, childhood maltreatment may relate to emotional sensitivity among those with secondary psychopathy. There is evidence that maltreatment adversely affects the development of the hypothalamic-pituitary-adrenocortical (HPA) system, which regulates emotional and particularly fearful responding (see Nelson & Carver, 1998). In turn, maltreated children are less likely to offer help to or show sadness or

concern towards a distressed peer, and are more likely to withdraw from or aggress against that peer (Klimes-Dougan & Kistner, 1990; Main & George, 1985). Genetically sensitive, longitudinal research is needed to explore how differences in emotional processing among primary and secondary variants of psychopathy may relate to temperamental and environmental factors.

Remarkably, neither variant processed distress significantly differently from nonpsychopathic offenders; comparison offenders fell mid-way between primary and secondary variants in their responsiveness to distressing stimuli. As previously mentioned, similar findings were reported by Bagley et al. (2009). This finding may reflect the considerable heterogeneity of offenders as a group. The spread of scores on the dot probe task was greater for general (nonpsychopathic) offenders (-310.67 to 382.37ms) than for primary (-251.75 to 113.96ms) and secondary psychopathy variants (-100.75 to 361.33ms), reflecting greater variability in the comparison group. There are at least two possible explanations for this. One explanation is that measures of psychopathy may lack precision in adolescent samples because of normative variations in the level of these traits across adolescents (Edens, Skeem, Cruise, & Cauffman, 2001). Alternatively, this heterogeneity in the general offending group may be tied to more divergent dispositional characteristics and contextual experiences across youth who enter the juvenile justice system. For example, youth high on anxiety (Vasey, et al., 1995; Watts & Weems, 2006), narcissism (Munoz, Kimonis, Strickleton, Frick, & Aucoin, 2010), and those with conduct problems/aggression without psychopathic traits (Kimonis, Frick, et al., 2006) tend to show heightened attention to negative emotional cues. Thus, the high prevalence of aggression, narcissism and anxiety disorders in incarcerated youth (Baumeister, Smart, & Boden, 1996; Karnik, et al., 2009) may account for the failure to find a significant difference from

secondary variants. In sum, although primary variants did not display an emotional *deficit* and secondary variants did not display an *enhancement* in attention to others' distress cues relative to the comparison group, our results do suggest that secondary variants were still more attentively engaged by images of others in distress than were primary variants.

The results of this study must be considered within the context of several study limitations. First, whereas prior research finds that anxiety is associated with hypervigilance towards threatening stimuli on the dot probe task (e.g., Vasey, et al., 1996), we were able to rule out the possibility that differences between psychopathy variants in emotional processing of distressing stimuli were entirely accounted for by differences in anxiety, as these two variables were not significantly correlated ($r = .10$). Second, the current study is limited to serious male juvenile offenders and findings may not generalize to girls, community samples, or to juvenile offenders with less serious histories of offending. That is, this study employed an adolescent offender sample housed in a secure facility, the majority of whom were sentenced for a violent committing offense. Indeed, average YPI scores were fairly high in this sample compared to a more heterogeneous sample of boys adjudicated of a felony offense (capped at 15%, Schubert, et al., 2004 vs. 4%, present study). Also, the sample consisted of predominately ethnic minority youth (72%), particularly those of Hispanic descent (47%), a population for which validation studies of the YPI have yet to be conducted. While this is an important strength, given the relative lack of research on CU traits and psychopathy in minority populations, it also limits the generalizability of findings to less ethnically heterogeneous populations. However, chi-square difference tests from multiple group analysis supported the covariance equivalence of the YPI and RCMAS measures across the two primary racial/ethnic groups (Black and Hispanic) included in this study. Third, as noted previously etiological inferences cannot be drawn from

this cross-sectional study. That is, emotional processing deficits may reflect temperamental differences in emotionality, or alternatively, may result from exposure to violent home and neighborhood environments that cause desensitization (see Kimonis, et al., 2008). For example, infants raised in severely impoverished social and emotional environments show a blunted pattern of emotional (i.e., cortisol) reactivity (Carlson & Earls, 1997; Gilles, Bernston, Zipf, & Gunnar, 2000), which may persist into adulthood (van der Vegt, van der Ende, Kirschbaum, Verhulst, & Tiemeier, 2009).

Implications for Future Research and Practice

Models to explain antisocial behavior need to explain the many differences between antisocial youth with and without high levels of CU traits. Our results further suggest that youth high on these traits can also be disaggregated into important subgroups that differ in their emotional stability and emotional processing. As suggested earlier, these results help shape etiologic hypotheses to address in future research. For example, our results suggest that future prospective studies could follow two groups of children: those with a fearless temperament and deficits in processing distressing emotional stimuli (i.e., primary psychopathy) and those with a disinhibited temperament who are exposed to high rates of trauma and abuse (i.e., secondary psychopathy). Fortunately, there are methods for assessing fearlessness and related temperamental dimensions in very young children (Fowles & Kochanska, 2000; Kagan, Reznik, & Snidman, 1988).

Such prospective studies could be very important for developing effective prevention programs for CU traits by identifying factors that may reduce the risk for problems in conscience development in youth with a fearful temperament. For example, Kochanska (Kochanska, 1997; Kochanska & Murray, 2000) proposed that the parent-child relationship, especially the

responsiveness towards each other, may be critical in the socialization of children with a fearless temperament. This aspect of parenting does not rely on punishment-related arousal for internalization and, instead focuses on the positive qualities of the parent-child relationship (Kochanska & Murray, 2000). In support of this proposal, attachment security was shown to be predictive of conscience development in temperamentally fearless children (Kochanska, 1995; 1997). Also, Cornell and Frick (2007) specifically tested several interactions between a fearless temperament and different dimensions of parenting in predicting scores on measures of guilt and empathy in young (age 3- 5 years) children. They reported two interactions, such that fearless children showed higher levels of guilt when parental consistency was high and when parents used authoritarian parenting (i.e., strong rule-oriented and obedience-oriented parenting). The authors interpreted these findings to suggest that fearless children require stronger and more consistent parenting to develop appropriate levels of guilt.

More immediately, our findings can shape hypotheses about targeted intervention for psychopathic juvenile offenders to evaluate in future research. At the outset, it is important to recognize that a growing body of research indicates that intensive treatment meaningfully reduces risk of violence and other criminal behavior for youth with psychopathic traits (for a review, see Skeem, Polaschek, & Manchak, 2009). However, it is possible that even greater gains may be made if treatment is targeted to the psychopathic variants identified in this study. For example, research suggests that cognitive-behavioral interventions may be most effective at treating internalizing problems (e.g., anger, anxiety, and depression) and related trauma histories (see Chaffin & Friedrich, 2004; Kaslow & Thompson, 1998; Ollendick & King, 1998) that distinguish secondary variants. For the low-anxious primary variant, recent research suggests that deficits in attention to others' distress cues can at least temporarily be corrected by focusing

youths' attention on the eye region (Dadds, et al., 2006) (see also Baskin-Sommers, Curtin, & Newman, 2011). Also, increasing the salience of others' distress cues has been found to attenuate laboratory-based aggression for youth scoring high on psychopathic traits (van Baardewijk, Stegge, Bushman, & Vermeiren, 2009). This group has also been shown to respond positively to rewards and this also can be capitalized on in treatment. For example, Hawes and Dadds (2005) reported that clinic-referred boys (ages 4 to 9) with conduct problems and CU traits were less responsive to a parenting intervention than were boys with conduct problems who were low on CU traits. However, children with and without CU traits seemed to respond equally well to the first part of the intervention that focused on teaching parents methods of using positive reinforcement to encourage prosocial behavior. In contrast, only the group without CU traits showed added improvement with the second part of the intervention that focused on teaching parents more effective discipline strategies. In summary, several promising interventions have emerged for youth with CU traits. These efforts are likely to be enhanced if they consider the heterogeneity among high CU youth and appropriately tailor treatment to their individual needs.

References

- Achenbach, T. M. (1991). *Integrative guide to the 1991 CBCL 14-18, YSR, and TRF profiles*. Burlington, VT: University of Vermont Department of Psychiatry.
- Andershed, H. A., Kerr, M., Stattin, H., & Levander, S. (2002). Psychopathic traits in non-referred youths: A new assessment tool. In E. Blauuw & L. Sheridan (Eds.), *Psychopaths: Current International Perspectives* (pp. 131-158). The Hague: Elsevier.
- Bagley, A. D., Abramowitz, C. S., & Kosson, D. S. (2009). Vocal affect recognition and psychopathy: Converging findings across traditional and cluster analytic approaches to assessing the construct. *Journal of Abnormal Psychology, 118*(2), 388-398.
- Banfield, J. D., & Raftery, A. E. (1993). Model-based Gaussian and non-Gaussian clustering. *Biometrics, 49*, 803–821.
- Bar-Haim, Y., Lamy, D., Pergamin, L., Bakermans-Kranenberg, M. J., & van Ijzendoorn, M. H. (2007). Threat-related attentional bias in anxious and nonanxious individuals: A meta-analytic study. *Psychological Bulletin, 133*, 1-24.
- Baskin-Sommers, A. R., Curtin, J. J., & Newman, J. P. (2011). Specifying the attentional selection that moderates the fearlessness of psychopathic offenders. *Psychological Science, 22*(2), 226-234.
- Baumeister, R. F., Smart, L., & Boden, J. M. (1996). Relation of threatened egotism to violence and aggression: The dark side of high self-esteem. *Psychological Review, 103*(1), 5-33.
- Blair, R. J. R. (1999). Responsiveness to distress cues in the child with psychopathic tendencies. *Personality and Individual Differences, 27*, 135-145.

- Blair, R. J. R., Colledge, E., Murray, L., & Mitchell, D. G. V. (2001). A selective impairment in the processing of sad and fearful expressions in children with psychopathic tendencies. *Journal of Abnormal Child Psychology, 29*(6), 491-498.
- Blair, R. J. R., Peschardt, K. S., Budhani, S., Mitchell, D. G. V., & Pine, D. S. (2006). The development of psychopathy. *Journal of Child Psychology and Psychiatry and Allied Disciplines, 47*(3), 262-275.
- Carlson, M., & Earls, F. (1997). Psychological and neuroendocrinological sequelae of early social deprivation in institutionalized children in Romania. *Annals of the New York Academy of Sciences, 807*, 419-428.
- Cauffman, E., Kimonis, E. R., Dmitrieva, J., & Monahan, K. (2009). A multi-method assessment of juvenile psychopathy: Comparing the predictive utility of the PCL:YV, YPI, and NEO PRI. *Psychological Assessment, 21*(4), 528-542.
- Chaffin, M., & Friedrich, B. (2004). Evidence-based treatments in child abuse and neglect. *Children and Youth Services Review, 26*, 1097-1113.
- Cleckley, H. (1941). *The mask of sanity: An attempt to reinterpret the so-called psychopathic personality* (1 ed.). Oxford, England.
- Cleckley, H. (1955). *The Mask of Sanity* (3rd ed.). St. Louis: C.V. Mosby.
- Cleckley, H. (1976). *The mask of sanity* (5th ed.). Saint Louis: C.V.: Mosby Co.
- Cooke, D. J., & Michie, C. (2001). Refining the construct of psychopathy: Towards a hierarchical model. *Psychological Assessment, 13*, 171-188.
- Cornell, A. H., & Frick, P. J. (2007). The moderating effects of parenting styles in the association between behavioral inhibition and parent-reported guilt and empathy in preschool children. *Journal of Clinical Child and Adolescent Psychology, 36*(3), 305-318.

- Dadds, M. R., El Masry, Y., Wimalaweera, S., & Guastella, A. J. (2008). Reduced eye gaze explains 'fear blindness' in childhood psychopathic traits. *Journal of the American Academy of Child & Adolescent Psychiatry, 47*(4), 455-463.
- Dadds, M. R., Perry, Y., Hawes, D. J., Merz, S., Riddell, A. C., Haines, D. J., et al. (2006). Look at the eyes: Fear recognition in child psychopathy. *British Journal of Psychiatry, 189*, 180-181.
- Dodge, K. A., & Pettit, G. S. (2003). A biopsychosocial model of the development of chronic conduct problems in adolescence. *Developmental Psychology, 39*, 349-371.
- Edens, J. F., Marcus, D. K., Lilienfeld, S. O., & Poythress, N. G. (2006). Psychopathic, Not Psychopath: Taxometric Evidence for the Dimensional Structure of Psychopathy. *Journal of Abnormal Psychology, 115*(1), 131-144.
- Edens, J. F., Skeem, J., Cruise, K., & Cauffman, E. (2001). The assessment of juvenile psychopathy and its association with violence: A critical review. *Behavioral Sciences & the Law, 19*, 53-80.
- Edens, J. F., Skopp, N. A., & Cahill, M. A. (2008). Psychopathic features moderate the relationship between harsh and inconsistent parental discipline and adolescent antisocial behavior. *Journal of Clinical Child and Adolescent Psychology, 37*, 472-476.
- Fisher, L., & Blair, R. J. R. (1998). Cognitive impairment and its relationship to psychopathic tendencies in children with emotional and behavioral difficulties. *Journal of Abnormal Child Psychology, 26*, 511-519.
- Fontaine, N. M. G., Rijdsdijk, F. V., McCrory, E. J. P., & Viding, E. (2010). Etiology of Different Developmental Trajectories of Callous-Unemotional Traits. *Journal of the American Academy of Child and Adolescent Psychiatry, 49*(7), 656 - 664.

- Forth, A. E., Kosson, D. S., & Hare, R. D. (2003). *The Psychopathy Checklist: Youth Version manual*. Toronto: Multi-Health Systems.
- Fowles, D. C., & Kochanska, G. (2000). Temperament as a moderator of pathways to conscience in children: The contribution of electrodermal activity. *Psychophysiology*, 37(6), 788-795.
- Fraley, C., & Raftery, A. E. (2002a). MCLUST: Software for model-based clustering, density estimation and discriminant analysis. Technical Report. Retrieved from <http://www.stat.washington.edu/tech.reports>
- Fraley, C., & Raftery, A. E. (2002b). Model-based clustering, discriminant analysis, and density estimation. *Journal of the American Statistical Association*, 380(2000).
- Fraley, C., & Raftery, A. E. (2008). The mclust02 Package, from <http://www.stat.washington.edu/mclust>
- Frick, P. J. (2006). Developmental pathways to conduct disorder. *Child Psychiatric Clinics of North America*, 15, 311-332.
- Frick, P. J. (2009). Extending the construct of psychopathy to youth: implications for understanding, diagnosing, and treating antisocial children and adolescents. *Canadian journal of psychiatry. Revue canadienne de psychiatrie*, 54(12), 803-812.
- Frick, P. J., Blair, R. J. R., & Castellanos, F. X. (in press). Callous-unemotional traits and developmental pathways to the disruptive behavior disorders. In P. H. Tolan & B. L. Leventhal (Eds.), *Advances in child and adolescent psychopathology Volume I: Disruptive Behavior Disorders*. New York: Springer.

- Frick, P. J., Cornell, A. H., Bodin, S. D., Dane, H. A., Barry, C. T., & Loney, B. R. (2003). Callous-unemotional traits and developmental pathways to severe conduct problems. *Developmental Psychology, 39*, 246-260.
- Frick, P. J., & Moffitt, T. E. (2010). *A Proposal to the DSM-V Childhood Disorders and the ADHD and Disruptive Behavior Disorders Work Groups to Include a Specifier to the Diagnosis of Conduct Disorder based on the Presence of Callous-Unemotional Traits*. Washington, DC: APA.
- Frick, P. J., & Morris, A. S. (2004). Temperament and developmental pathways to conduct problems. *Journal of Clinical Child and Adolescent Psychology, 33*, 54-68.
- Frick, P. J., Stickle, T. R., Dandreaux, D. M., Farrell, J. M., & Kimonis, E. R. (2005). Callous-Unemotional Traits in Predicting the Severity and Stability of Conduct Problems and Delinquency. *Journal of Abnormal Child Psychology, 33*(4), 471-487.
- Frick, P. J., & Viding, E. M. (2009). Antisocial behavior from a developmental psychopathology perspective. *Development and Psychopathology, 21*, 1111-1131.
- Gil-Rivas, V. (2003). *Parental Contributions to Adolescents' Psychological Adjustment Following the Terrorist Attacks of September 11, 2001*. Unpublished doctoral dissertation. University of California, Irvine.
- Gilles, E. E., Bernston, G. G., Zipf, W. B., & Gunnar, M. R. (2000). *Neglect is associated with a blunting of the behavioral and biological stress responses in human infants*. Paper presented at the International Conference of Infant Studies, Brighton, England.
- Guay, J. P., Ruscio, J., Knight, R. A., & Hare, R. D. (2007). A Taxometric Analysis of the Latent Structure of Psychopathy: Evidence for Dimensionality. *Journal of Abnormal Psychology, 116*(4), 701-716.

- Hare, R. D. (2003). *Hare Psychopathy Checklist-Revised (PCL-R): 2nd Edition, Technical Manual*. Canada: Multi-Health Systems Inc.
- Hare, R. D., & Neumann, C. (2010). The role of antisociality in the psychopathy construct: Comment on Skeem and Cooke. *Psychological Assessment, 22*(2), 446-454.
- Harris, G. T., Rice, M. E., & Quinsey, V. L. (1994). Psychopathy as a taxon: Evidence that psychopaths are a discrete class. *Journal of Consulting and Clinical Psychology, 62*(2), 387-397.
- Hart, S. D., Cox, D., & Hare, R. D. (1995). *Manual for the Psychopathy Checklist: Screening Version (PCL: SV)*. Toronto: Multi-Health Systems.
- Hawes, D. J., & Dadds, M. R. (2005). The treatment of conduct problems in children with callous-unemotional traits. *Journal of Consulting and Clinical Psychology, 73*(4), 737-741.
- Hicks, B. M., Markon, K., Patrick, C. J., Krueger, R., & Newman, J. P. (2004). Identifying psychopathy subtypes on the basis of personality structure. *Psychological Assessment, 16*, 276-288.
- Hicks, B. M., Vaidyanathan, U., & Patrick, C. J. (2010). Validating female psychopathy subtypes: Differences in personality, antisocial and violent behavior, substance abuse, trauma, and mental health. *Personality Disorders: Theory, Research, and Treatment, 1*(1), 38-57.
- Hipwell, A. E., Pardini, D. A., Loeber, R., Sembower, M., Keenan, K., & Stouthamer-Loeber, M. (2007). Callous-unemotional behaviors in young girls: Shared and unique effects relative to conduct problems. *Journal of Clinical Child and Adolescent Psychology, 36*(3), 293-304.

- Hoffman, M. L. (1982). Development of prosocial motivation: Empathy and guilt. In N. Eisenberg (Ed.), *The development of prosocial behavior* (pp. 281-313). San Diego, CA: Academic Press.
- Insightful Corporation. (1988-2005). S-PLUS version 7.0 for Windows. Seattle, Washington: Insightful Corporation.
- Jones, A. P., Laurens, K. L., Herba, C., Barker, G., & Viding, E. (2009). Amygdala hypoactivity to fearful faces in boys with conduct problems and callous-unemotional traits. *American Journal of Psychiatry, 166*, 95-102.
- Kagan, J., Reznik, J. S., & Snidman, N. (1988). Biological bases of childhood shyness. *Science, 240*, 167-171.
- Kahn, R. E., Frick, P. J., Youngstrom, E., Findling, R. L., & Youngstrom, J. K. (in press). The effects of including a callous-unemotional specifier for the diagnosis of conduct disorder. *Journal of Child Psychology and Psychiatry*.
- Karnik, N. S., Soller, M., Redlich, A., Silverman, M., Kraemer, H. C., Haapanen, R., et al. (2009). Prevalence of and gender differences in psychiatric disorders among juvenile delinquents incarcerated for nine months. *Psychiatric Services, 60*(6), 838-841.
- Karpman, B. (1941). On the need of separating psychopathy into two distinct clinical types: The symptomatic and the idiopathic. *Journal of Criminology and Psychopathology, 3*, 112-137.
- Karpman, B. (1948a). Conscience in the psychopath: Another version. *American Journal of Orthopsychiatry, 18*, 455-491.
- Karpman, B. (1948b). The myth of the psychopathic personality. *American Journal of Psychiatry, 104*, 523-534.

- Kaslow, N. J., & Thompson, M. P. (1998). Applying the criteria for empirically supported treatments to studies of psychosocial interventions for child and adolescent depression. *Journal of Clinical Child Psychology, 27*(2), 146-155.
- Kimonis, E. R., Cauffman, E., & Goldweber, A. (2009, April). *Emotional Deficits in Youth with Callous-Unemotional Traits: Consistency Across Ethnicity and Gender?* Paper presented at the Society for the Scientific Study of Psychopathy conference, New Orleans, LA.
- Kimonis, E. R., Frick, P. J., Fazekas, H., & Loney, B. R. (2006). Psychopathy, aggression, and the processing of emotional stimuli in non-referred girls and boys. *Behavioral Sciences & the Law. Special Issue: Gender and Psychopathy Volume 2, 24*(1), 21-37.
- Kimonis, E. R., Frick, P. J., Muñoz, L. C., & Aucoin, K. J. (2008). Callous-Unemotional Traits and the Emotional Processing of Distress Cues in Detained Boys: Testing the Moderating Role of Aggression, Exposure to Community Violence, and Histories of Abuse. *Development and Psychopathology, 20*, 569-589.
- Kimonis, E. R., Skeem, J., Cauffman, E., & Dmitrieva, J. (2011). Are secondary variants of 'juvenile psychopathy' more reactively violent and less psychosocially mature than primary variants? *Law and Human Behavior, 35*(5), 381-391. doi: 10.1007/s10979-010-9243-3
- Kimonis, E. R., Skeem, J. L., & Cauffman, E. (2006). *Psychopathy-like juvenile offenders: Are secondary variants less mature and stable than primary variants?* Paper presented at the Society for Research on Adolescence, San Francisco, CA.
- Klimes-Dougan, B., & Kistner, J. (1990). Physically abused preschoolers' responses to peers' distress. *Developmental Psychology, 26*, 599-602.

- Kochanska, G. (1993). Toward a synthesis of parental socialization and child temperament in early development of conscience. *Child Development, 64*, 325-347.
- Kochanska, G. (1995). Children's temperament, mothers' discipline, and security of attachment: Multiple pathways to emerging internalization. *Child Development, 66*, 597-615.
- Kochanska, G. (1997). Multiple pathways to conscience for children with different temperaments: From toddlerhood to age 5. *Developmental Psychology, 33*, 228-240.
- Kochanska, G., & Murray, K. T. (2000). Mother-child mutually responsive orientation and conscience development: From toddler to early school age. *Child Development, 71*, 417-431.
- Kruh, I. P., Frick, P. J., & Clements, C. B. (2005). Historical and personality correlates to the violence patterns of juveniles tried as adults. *Criminal Justice and Behavior, 32*, 69-96.
- Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics, 33*, 159-174.
- Lang, P. J., Bradley, M. M., & Cuthbert, B. N. (1997). *International affective picture system (IAPS)*: NIMH Center for the Study of Emotions and Attention.
- Lee, Z., Salekin, R., & Iselin, A. R. (2010). Psychopathic Traits in Youth: Is There Evidence for Primary and Secondary Subtypes? *Journal of Abnormal Child Psychology, 38*, 381-393.
doi: 10.1007/s10802-009-9372-7
- Legerstee, J. S., Tullen, J. H. M., Kallen, V. L., Dieleman, G. C., Treffers, P. D. A., Verhulst, F. C., et al. (2009). Threat-related selective attention predicts treatment success in childhood anxiety disorders. *Journal of the American Academy of Child & Adolescent Psychiatry, 48*(2), 196-205.

- Lilienfeld, S. (1994). Conceptual problems in the assessment of psychopathy. *Clinical Psychology Review, 14*, 17-38.
- Lilienfeld, S. (1998). Methodological advances and developments in the assessment of psychopathy. *Behaviour Research and Therapy, 36*, 99-125.
- Loney, B. R., Butler, M. A., Lima, E. N., Counts, C. A., & Eckel, L. A. (2006). The relation between salivary cortisol, callous-unemotional traits, and conduct problems in an adolescent non-referred sample. *Journal of Child Psychology and Psychiatry, 47*(1), 30-36.
- Lykken, D. T. (1957). A study of anxiety in the sociopathic personality. *Journal of Abnormal and Social Psychology, 55*, 6-10.
- Lykken, D. T. (1995). *The antisocial personalities*. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc.
- MacLeod, C., & Mathews, A. (1988). Anxiety and the allocation of attention to threat. *Quarterly Journal of Experimental Psychology A. Human Experimental Psychology, 40*(4-A), 653-670.
- MacLeod, C., Mathews, A., & Tata, P. (1986). Attentional bias in emotional disorders. *Journal of Abnormal Psychology, 95*(1), 15-20.
- Main, M., & George, C. (1985). Responses of abused and disadvantaged toddlers to distress inagemates: A study in the day care setting. *Developmental Psychology, 21*, 407-412.
- Marcus, D. K., Lilienfeld, S. O., Edens, J. F., & Poythress, N. G. (2006). Is antisocial personality disorder continuous or categorical? A taxometric analysis. *Psychological Medicine, 36*(11), 1571-1581.

Marsh, A. A., Finger, E. C., Mitchell, D. G., Reid, M. E., Sims, C., Kosson, D. S., et al. (2008).

Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behavior disorders. *American Journal of Psychiatry*, *165*, 712-720.

Moffitt, T. E. (1993). Adolescence-limited and life-course persistent antisocial behavior: A developmental taxonomy. *Psychological Review*, *100*, 674-701.

Munoz, L. C., Kimonis, E. R., Strickleton, C., Frick, P. J., & Aucoin, K. (2010). *Heightened Emotional Reactivity and Proactive Aggression in Youth with Narcissistic Traits*. Paper presented at the American Psychology-Law Society conference, Vancouver, Canada.

Muris, P., Merckelbach, H., Ollendick, T., King, N., & Bogie, N. (2002). Three traditional and three new childhood anxiety questionnaires: Their reliability and validity in a normal adolescent sample. *Behaviour Research and Therapy*, *40*(7), 753-772.

Murrie, D. C., Marcus, D. K., Douglas, K., Lee, Z., Salekin, R. T., & Vincent, G. M. (2007). Youth with psychopathy features are not a discrete class: a taxometric analysis. *Journal of Child Psychology and Psychiatry*, *48*(7), 714-723.

Muthén, L. K., & Muthén, B. O. (2007). *Mplus user's guide (5th edition)* (5th ed.). Los Angeles, CA: Muthén & Muthén.

Nelson, C. A., & Carver, L. J. (1998). The effects of stress and trauma on brain and memory: A view from developmental cognitive neuroscience. *Development and Psychopathology*, *10*(4), 793-809.

Newman, J. P., Patterson, C. M., Howland, E. W., & Nichols, S. L. (1990). Passive avoidance in psychopaths: The effects of reward. *Personality and Individual Differences*, *11*, 1101-1114.

- Newman, J. P., Schmitt, W., & Voss, W. (1997). The impact of motivational neutral cues on psychopaths: Assessing the generality of the response modulation hypothesis. *Journal of Abnormal Psychology, 106*, 563-575.
- Newman, J. P., & Schmitt, W. A. (1998). Passive avoidance in psychopathic offenders: A replication and extension. *Journal of Abnormal Psychology, 107* (3), 527-532.
- Novaco, R. W. (2003). *The Novaco Anger Scale and Provocation Inventory (NAS-PI)*. Los Angeles, CA: Western Psychological Services.
- Ollendick, T., & King, N. (1998). Empirically supported treatments for children with phobic and anxiety disorders: Current status. *Journal of Clinical Child Psychology, 27*(2), 156-167.
- Oxford, M., Cavell, T. A., & Hughes, J. N. (2003). Callous-unemotional traits moderate the relation between ineffective parenting and child externalizing problems: A partial replication and extension. *Journal of Clinical Child and Adolescent Psychology, 32*, 577-585.
- Pardini, D. A., Lochman, J. E., & Frick, P. J. (2003). Callous/unemotional traits and social-cognitive processes in adjudicated youths. *Journal of the American Academy of Child & Adolescent Psychiatry, 42*(3), 364-371.
- Patrick, C. J., Fowles, D. C., & Krueger, R. (2009). Triarchic conceptualization of psychopathy: Developmental origins of disinhibition, boldness, and meanness. *Development and Psychopathology, 21*, 913-938.
- Pollak, S. D., & Kistler, D. J. (2002). Early experience is associated with the development of categorical representations for facial expressions of emotion. *Proceedings of the National Academy of Sciences, 99*, 9072-9076.

- Pollak, S. D., Klorman, R., Thatcher, J. E., & Cicchetti, D. (2001). P3b reflects maltreated children's reactions to facial displays of emotion. *Psychophysiology*, *38*, 267-274.
- Pollak, S. D., & Sinha, P. (2002). Effects of early experience on children's recognition of facial displays of emotion. *Developmental Psychology*, *38*, 784-791.
- Pollak, S. D., & Tolley-Schell, S. A. (2003). Selective attention to facial emotion in physically abused children. *Journal of Abnormal Psychology*, *112*, 323-338.
- Poythress, N. G., Edens, J. F., Skeem, J. L., Lilienfeld, S. O., Douglas, K. S., Frick, P. J., et al. (2010). Identifying Subtypes among Offenders with Antisocial Personality Disorder: A Cluster-Analytic Study. *Journal of Abnormal Psychology*.
- Poythress, N. G., & Skeem, J. L. (2005). Disaggregating psychopathy: Where and how to look for variants. In C. J. Patrick (Ed.), *Handbook of Psychopathy* (pp. 172-192). New York, NY: Guilford Press.
- Radloff, L. (1977). The CES-D scale: a self-report depression scale for research in the general population. *Applied Psychological Measurement*, *1*, 385-401.
- Reynolds, C. R., & Richmond, B. O. (1985). *Revised Children's Manifest Anxiety Scale*. Los Angeles, CA: Western Psychological Services.
- Reynolds, C. R., & Richmond, B. O. (2000). *Revised Children's Manifest Anxiety Scale: Manual*. Los Angeles, CA: Western Psychological Services.
- Rowe, R., Maughan, B., Moran, P., Ford, T., Briskman, J., & Goodman, R. (2010). The role of callous and unemotional traits in the diagnosis of conduct disorder. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, *51*(6), 688-695.
- Schippell, P. L., Vasey, M. W., Cravens-Brown, L. M., & Bretveld, R. A. (2003). Suppressed attention to rejection, ridicule, and failure cues: A unique correlate of reactive but not

- proactive aggression in youth. *Journal of Clinical Child and Adolescent Psychology*, 32, 40-55.
- Schubert, C., Mulvey, E., Steinberg, L., Cauffman, E., Losoya, S., Hecker, T., et al. (2004). Operational lessons from the pathways to desistance project. *Journal of Youth Violence and Juvenile Justice*, 2, 237-255.
- Skeem, J. L., & Cooke, D. J. (2010a). Is criminal behavior a central component of psychopathy? Conceptual directions for resolving the debate. *Psychological Assessment*, 22(2), 433-445.
- Skeem, J. L., & Cooke, D. J. (2010b). One measure does not a construct make: Directions toward reinvigorating psychopathy research—reply to Hare and Neumann (2010). *Psychological Assessment*, 22(2), 455-459.
- Skeem, J. L., Johansson, P., Andershed, H. A., Kerr, M., & Louden, J. E. (2007). Two subtypes of psychopathic violent offenders that parallel primary and secondary variants. *Journal of Abnormal Psychology*, 116(2), 395-409.
- Skeem, J. L., Polaschek, D. L. L., & Manchak, S. (2009). Appropriate treatment works, but how?: Rehabilitating general, psychopathic, and high-risk offenders. In J. L. Skeem, K. S. Douglas & S. O. Lilienfeld (Eds.), *Psychological science in the courtroom: Consensus and controversy* (pp. 358-384). New York, NY: Guilford Press.
- Skeem, J. L., Poythress, N. G., Edens, J. F., Lilienfeld, S. O., & Cale, E. M. (2003). Psychopathic personality or personalities? Exploring potential variants of psychopathy and their implications for risk assessment. *Aggression and Violent Behavior*, 8(5), 513-546.

Snyder, H. N., & Sickmund, M. (2006). *Juvenile Offenders and Victims: 2006 National Report*.

Washington, DC: U.S. Department of Justice, Office of Justice Programs, Office of Juvenile Justice and Delinquency Prevention.

Tatar, J. R., Cauffman, E., Kimonis, E. R., & Skeem, J. L. (in press). Victimization history and post-traumatic stress: An analysis of psychopathy variants in male juvenile offenders.

Journal of Child and Adolescent Trauma.

van Baardewijk, Y., Stegge, H., Bushman, B. J., & Vermeiren, R. (2009). Psychopathic traits, victim distress and aggression in children. *Journal of Child Psychology and Psychiatry*, *50*(6), 718-725. doi: 10.1111/j.1469-7610.2008.02023.x

van der Vegt, E. J. M., van der Ende, J., Kirschbaum, C., Verhulst, F. C., & Tiemeier, H. (2009).

Early neglect and abuse predict diurnal cortisol patterns in adults: A study of international adoptees. *Psychoneuroendocrinology*, *34*(5), 660-669. doi: 10.1016/j.psyneuen.2008.11.004

Vasey, M. W., Daleidon, E. L., Williams, L. L., & Brown, L. M. (1995). Biased attention in childhood anxiety disorders: A preliminary study. *Journal of Abnormal Child Psychology*, *23*, 267-279.

Vasey, M. W., El-Hag, N., & Daleidon, E. L. (1996). Anxiety and the processing of emotionally threatening stimuli: Distinctive patterns of selective attention among high and low-test-anxious children. *Child Development*, *67*(1173-1185).

Vasey, M. W., Kotov, R., Frick, P. J., & Loney, B. R. (2005). The latent structure of psychopathy in youth: A taxometric investigation. *Journal of Abnormal Child Psychology*, *33*(4), 411-429.

- Vassileva, J., Kosson, D. S., Abramowitz, C., & Conrod, P. (2005). Psychopathy versus psychopathies in classifying criminal offenders. *Legal & Criminological Psychology, 10*, 27-43.
- Vaughn, M. G., Edens, J., Howard, M. O., & Smith, S. T. (2009). An investigation of primary and secondary psychopathy in a statewide sample of incarcerated youth. *Youth Violence and Juvenile Justice, 7*(3), 172-188.
- Viding, E., Blair, R. J. R., Moffitt, T. E., & Plomin, R. (2005). Evidence for a substantial genetic risk for psychopathic traits in 7-year-olds. *Journal of Child Psychology and Psychiatry, 46*(6), 592-597.
- Viding, E., Jones, A. P., Frick, P. J., Moffitt, T. E., & Plomin, R. (2008). Heritability of antisocial behaviour at 9: Do callous-unemotional traits matter? *Developmental Science, 11*(1), 17-22.
- Watts, S. E., & Weems, C. F. (2006). Associations among selective attention, memory bias, cognitive errors and symptoms of anxiety in youth. *Journal of Abnormal Child Psychology, 34*, 841-852. doi: 10.1007/s10802-006-9066-3
- Wootton, J. M., Frick, P. J., Shelton, K. K., & Silverthorn, P. (1997). Ineffective parenting and childhood conduct problems: The moderating role of callous-unemotional traits. *Journal of Consulting and Clinical Psychology, 65*, 301-308.
- Young, S. K., Fox, N. A., & Zahn-Waxler, C. (1999). The relations between temperament and empathy in 2-year-olds. *Developmental Psychology, 35*(5), 1189-1179.

Footnotes

¹Our use of the term *variants* as opposed to *subtypes* is for the purpose of identifying prototypes rather than discrete categories of youth, consistent with the most compelling data suggesting that psychopathic traits are dimensional rather than a taxon (adults, Edens, et al., 2006; Guay, Ruscio, Knight, & Hare, 2007; Lilienfeld, 1994; 1998; youth, Murrie, et al., 2007) (cf. adults, Harris, et al., 1994; youth, Vasey, et al., 2005).

²The accuracy of cluster membership as determined by MBC analysis was further verified by a second cluster analysis conducted using mixture modeling in Mplus (Muthén & Muthén, 2007). Again, a two-group solution provided the best fit to the data (BIC = 5226.8). An interrater reliability analysis using the Kappa statistic was performed to determine consistency among the two cluster solutions, resulting in Kappa = 0.98 ($p < 0.001$). As a rule of thumb, Kappa values ranging between 0.40 and 0.59 are considered moderate, 0.60 to 0.79 substantial, and 0.80 and above outstanding (Landis & Koch, 1977). Compared with the results of the MBC analysis, the resulting mixture modeling groups revealed a somewhat larger secondary group ($n = 47$) and a somewhat smaller primary group ($n = 118$). Most notably, however, the original 43 secondary variants identified by MBC were also assigned to the secondary group identified using mixture modeling. To maintain consistency with prior research, groups formed using MBC were used for analyses.

³Prior research with this same sample documented that secondary psychopathy variants report a greater history of stressful life events (Tatar, et al., in press). To address the possibility that parent-child interactions and attachment associated with maltreatment are of greater importance etiologically to the development of psychopathy subtypes than exposure to other stressful life events, which are found to result in dysregulation of the HPA axis, we ran Poisson

regression analyses with subtype membership predicting maltreatment controlling for stressful life events, and vice versa. Results indicated that subtype membership significantly predicted maltreatment after controlling for stressful life events ($B = -.22$, Wald $\chi^2 = 4.05$, $p < .05$); however, stressful life events were not significantly predicted after controlling for maltreatment ($B = -.05$, Wald $\chi^2 = .81$, $p = n.s$). Stressful life events were measured using a count of 25 items assessing a variety of stressors (e.g., serious accident, death of close family members, witnessed violence [not domestic], tragedy or natural disaster and other such traumatic events)—not including maltreatment experiences—from the Stressful Life Events scale (Gil-Rivas, 2003). These results support the possibility that maltreatment may be more relevant to the development of secondary psychopathy than stressful life events more generally.

Table 1

Characteristics of the Total Sample, the Non-psychopathic Comparison Sample, and the Clustering Sample (YPI > 121.5)

Variable	Total sample (N=373)	Non- psychopathic comparison sample (n=208)	Clustering sample (YPI > 121.5; n=165)	Sig.
Mean Age (baseline)	16.42 (.79)	16.39 (.84)	16.47 (.74)	n.s.
Ethnicity				
White	6%	8%	5%	n.s.
Black	29%	27%	31%	
Hispanic	53%	44%	50%	
Current Offense				person: χ^2
Person	69%	75%	63%	(1,372)=5.92,
Property	12%	6%	19%	$p < .05$; property:
Drug offense	4%	4%	3%	$\chi^2(1,372)=14.08$,
				$p < .001$; drug: n.s.
Total YPI Mean (SD)	118.31 (20.48)	103.53 (13.87) ^a	135.43 (11.66) ^b	$t(1,353) = -23.57$;
				$p < .001$
Total Anxiety Mean (SD)	6.87 (5.19)	6.07 (5.24) ^a	7.80 (5.00) ^b	$t(1,354) = -3.17$;
				$p < .01$
Facilitation to Distress (ms)	-19.19 (56.30)	-17.87 (50.30)	-20.82 (63.08)	n.s.
Facilitation to Positive (ms)	-2.78 (71.31)	-6.09 (49.84)	1.27 (90.99)	n.s.

Note: YPI = Youth Psychopathic Traits Inventory. Facilitation scores are based on performance on the emotional pictures dot-probe task. Superscripts indicate a significant difference between the comparison subsample and the clustering sample.